



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

$$= \frac{1}{a^2 \sqrt{2}} \left[\frac{x_1/(x^2 + a^2)}{2} + \frac{a^2 \log[x + \sqrt{(x^2 + a^2)}]}{2} \right]_0^a = \frac{1}{2} \left[1 + \frac{\log(\sqrt{2} + 1)}{\sqrt{2}} \right].$$

O. W. Anthony gets as a result $\frac{2}{\pi}$. Professor MATZ furnished two solutions.

22. Proposed by ALTON L. SMITH, Instructor in Drawing, Polytechnic Institute, Worcester, Massachusetts.

In a series of counts of the votes on a legislative act relative to the city of Worcester, the following results were obtained:

	YES	NO
1st count	5566	5511
2nd "	5519	5558
3rd "	5546	5517
4th "	5512	5551
5th "	5512	5541

What is the probability that the last count (the 5th) is correct?

Solution by F. P. MATZ, M. Sc., Ph. D., Professor of Mathematics and Astronomy in New Windsor, College, New Windsor, Maryland.

I. Since the counts taken independently must be either correct or incorrect the probability that the fifth count is *incorrect* is

$P'_5 = \frac{1}{2} \times (\frac{1}{2} + \frac{1}{2} + \frac{1}{2} + \frac{1}{2} + \frac{1}{2}) = \frac{1}{2}$. Hence the probability that the fifth count is *correct* is $P_5 = 1 - P'_5 = \frac{1}{2}$.

II. According to *The Law of Experience*, the second count should show a *greater* probability as to correctness and a *smaller* probability as to incorrectness than the first count shows; that is, the probability as to the correctness of the fifth count should be *greater* than is the *similar* probability with respect to any other count *lower* than the fifth. This Law, according to the notation adopted, gives $P'_1 = \frac{1}{2}, P'_2 = \frac{1}{4}, \dots, P'_5 = \frac{1}{32} = (\frac{1}{2})^5$. Hence $P_5 = 1 - (\frac{1}{2})^5 = \frac{31}{32}$, which is the probability that the fifth count is correct.

PROBLEMS.

29. Proposed by JOHN DOLMAN, Jr., Philadelphia, Pennsylvania.

Neglecting perturbations, what is the average distance of the earth from the sun?

30. Proposed by F. P. MATZ, M. A., M. Sc., Ph. D., Professor of Mathematics and Astronomy in New Windsor College, New Windsor, Maryland.

Find the average area of all the triangles that can be inscribed in a given circle.

